

ARITHMETIC PROPERTIES OF q -FIBONACCI NUMBERS AND q -PELL NUMBERS

HAO PAN

ABSTRACT. We investigate some arithmetic properties of the q -Fibonacci numbers and the q -Pell numbers.

1. INTRODUCTION

The Fibonacci numbers F_n are given by

$$F_0 = 0, F_1 = 1 \text{ and } F_n = 2F_{n-1} + F_{n-2} \text{ for } n \geq 2.$$

For any odd prime p , it is well-known (cf. [7]) that

$$F_p \equiv \left(\frac{5}{p}\right) \pmod{p}, \quad (1.1)$$

$$F_{p+1} \equiv \frac{1}{2} \left(1 + \left(\frac{5}{p}\right)\right) \pmod{p} \quad (1.2)$$

and

$$F_{p-1} \equiv \frac{1}{2} \left(1 - \left(\frac{5}{p}\right)\right) \pmod{p}, \quad (1.3)$$

where $\left(\frac{\cdot}{p}\right)$ denotes the Legendre symbol. Indeed, we have

$$F_p = \frac{(1 + \sqrt{5})^p - (1 - \sqrt{5})^p}{2^p \sqrt{5}} \equiv \frac{(1 + (\sqrt{5})^p) - (1 - (\sqrt{5})^p)}{2\sqrt{5}} = 5^{(p-1)/2} \pmod{p}.$$

For more results on the congruences involving the Fibonacci numbers, the readers may refer to [10], [11] and [12].

On the other hand, a sequence of polynomials $\mathcal{F}_n(q)$ was firstly introduced by Schur (cf. [9]):

$$\mathcal{F}_n(q) = \begin{cases} 0 & \text{if } n = 0, \\ 1 & \text{if } n = 1, \\ \mathcal{F}_{n-1}(q) + q^{n-2} \mathcal{F}_{n-2}(q) & \text{if } n \geq 2. \end{cases}$$

1991 *Mathematics Subject Classification.* Primary 11B39; Secondary 05A30, 11B65, 11A07.

Key words and phrases. q -Fibonacci number, q -Pell number, Legendre symbol.

Also Schur considered another sequence $\widehat{\mathcal{F}}_n(q)$, which is given by

$$\widehat{\mathcal{F}}_n(q) = \begin{cases} 0 & \text{if } n = 0, \\ 1 & \text{if } n = 1, \\ \widehat{\mathcal{F}}_{n-1}(q) + q^{n-1}\widehat{\mathcal{F}}_{n-2}(q) & \text{if } n \geq 2. \end{cases}$$

Obviously both $\mathcal{F}_n(q)$ and $\widehat{\mathcal{F}}_n(q)$ are the q -analogues of the Fibonacci numbers. The sequences $\mathcal{F}_n(q)$ and $\widehat{\mathcal{F}}_n(q)$ have been investigated in several papers (e.g., see [2], [3], [4], [5] and [6]). However, seemingly there are no simple expressions for $\mathcal{F}_n(q)$ and $\widehat{\mathcal{F}}_n(q)$.

Now we can give the q -analogues of (1.1), (1.2) and (1.3). Suppose that $p \neq 5$ is an odd prime. Let α_p be the integer such that $1 \leq \alpha_p \leq 4$ and $\alpha_p p \equiv 1 \pmod{5}$. As usual we set

$$[n]_q = \frac{1 - q^n}{1 - q} = 1 + q + \cdots + q^{n-1}$$

for any non-negative integer n .

Theorem 1.1.

$$\mathcal{F}_{p+1} \equiv \frac{1}{2} \left(1 + \left(\frac{5}{p} \right) \right) \pmod{[p]_q} \quad (1.4)$$

and

$$\mathcal{F}_p \equiv \left(\frac{5}{p} \right) q^{((5-\alpha_p)p+1)/5} \pmod{[p]_q}. \quad (1.5)$$

Theorem 1.2.

$$\widehat{\mathcal{F}}_{p-1} \equiv \frac{1}{2} \left(1 - \left(\frac{5}{p} \right) \right) \pmod{[p]_q} \quad (1.6)$$

and

$$\widehat{\mathcal{F}}_p \equiv \left(\frac{5}{p} \right) q^{(\alpha_p p - 1)/5} \pmod{[p]_q}. \quad (1.7)$$

The Pell numbers P_n are given by

$$P_0 = 0, \quad P_1 = 1 \quad \text{and} \quad P_n = 2P_{n-1} + P_{n-2} \quad \text{for } n \geq 2.$$

It is easy to check that

$$P_n = \frac{(1 + \sqrt{2})^n - (1 - \sqrt{2})^n}{2\sqrt{2}}.$$

Hence for odd prime p , we have

$$P_p = \frac{(1 + \sqrt{2})^p - (1 - \sqrt{2})^p}{2\sqrt{2}} \equiv \frac{2(\sqrt{2})^p}{2\sqrt{2}} = 2^{(p-1)/2} \equiv \left(\frac{2}{p} \right) \pmod{p}. \quad (1.8)$$

Define the q -Pell numbers $\mathcal{P}_n(q)$ and $\widehat{\mathcal{P}}_n(q)$ by

$$\mathcal{P}_n(q) = \begin{cases} 0 & \text{if } n = 0, \\ 1 & \text{if } n = 1, \\ (1 + q^{n-1})\mathcal{P}_{n-1}(q) + q^{n-2}\mathcal{P}_{n-2}(q) & \text{if } n \geq 2, \end{cases}$$

and

$$\widehat{\mathcal{P}}_n(q) = \begin{cases} 0 & \text{if } n = 0, \\ 1 & \text{if } n = 1, \\ (1 + q^{n-1})\widehat{\mathcal{P}}_{n-1}(q) + q^{n-1}\widehat{\mathcal{P}}_{n-2}(q) & \text{if } n \geq 2. \end{cases}$$

Theorem 1.3. *Let p be an odd prime. Then*

$$q^{(p^2-1)/8}\mathcal{P}_p(q) \equiv \left(\frac{2}{p}\right) \pmod{[p]_q} \quad (1.9)$$

and

$$\widehat{\mathcal{P}}_p(q) \equiv \left(\frac{2}{p}\right) q^{(p^2-1)/8} \pmod{[p]_q}. \quad (1.10)$$

Furthermore, we have

$$\mathcal{P}_{p+1}(q) - \mathcal{P}_p(q) \equiv \widehat{\mathcal{P}}_{p+1}(q) - \widehat{\mathcal{P}}_p(q) \equiv 1 \pmod{[p]_q}. \quad (1.11)$$

Since $\left(\frac{2}{p}\right) = (-1)^{(p^2-1)/8}$, (1.9) and (1.10) can be respectively rewritten as

$$(-q)^{(p^2-1)/8}\mathcal{P}_p(q) \equiv 1 \pmod{[p]_q}$$

and

$$\widehat{\mathcal{P}}_p(q) \equiv (-q)^{(p^2-1)/8} \pmod{[p]_q}.$$

The proofs of Theorems 1.1, 1.2 and 1.3 will be given in Sections 2 and 3.

2. PROOFS OF THEOREMS 1.1 AND 1.2

For any $n, m \in \mathbb{Z}$, the q -binomial coefficient $\begin{bmatrix} n \\ m \end{bmatrix}_q$ is given by

$$\begin{bmatrix} n \\ m \end{bmatrix}_q = \frac{(1 - q^n)(1 - q^{n-1}) \cdots (1 - q^{n-m+1})}{(1 - q^m)(1 - q^{m-1}) \cdots (1 - q)}$$

when $m \geq 0$, and let $\begin{bmatrix} n \\ m \end{bmatrix}_q = 0$ if $m < 0$. Obviously $\begin{bmatrix} n \\ m \end{bmatrix}_q$ is a polynomial in the variable q with integral coefficients since the q -binomial coefficients satisfy the recurrence relation

$$\begin{bmatrix} n+1 \\ m \end{bmatrix}_q = q^m \begin{bmatrix} n \\ m \end{bmatrix}_q + \begin{bmatrix} n \\ m-1 \end{bmatrix}_q.$$

Let $\lfloor x \rfloor$ denotes the greatest integer not exceeding x . Then for any non-negative integer n , we have

$$\begin{aligned}\mathcal{F}_{n+1}(q) &= \sum_{0 \leq 2j \leq n} q^{j^2} \begin{bmatrix} n-j \\ j \end{bmatrix}_q \\ &= \sum_{j=-\infty}^{\infty} (-1)^j q^{j(5j+1)/2} \begin{bmatrix} n \\ \lfloor (n-5j)/2 \rfloor \end{bmatrix}_q\end{aligned}\quad (2.1)$$

and

$$\begin{aligned}\widehat{\mathcal{F}}_{n+1}(q) &= \sum_{0 \leq 2j \leq n} q^{j^2+j} \begin{bmatrix} n-j \\ j \end{bmatrix}_q \\ &= \sum_{j=-\infty}^{\infty} (-1)^j q^{j(5j-3)/2} \begin{bmatrix} n+1 \\ \lfloor (n+1-5j)/2 \rfloor + 1 \end{bmatrix}_q.\end{aligned}\quad (2.2)$$

We mention that (2.1) and (2.2) can be considered as the finite forms of the first and the second of the Rogers-Ramanujan identities respectively (the full proofs of (2.1) and (2.2) can be found in [1]).

In this section we assume that $p \neq 5$ is an odd prime.

Lemma 2.1. *Let*

$$L(j) = \frac{j(5j+1)}{2} - \begin{pmatrix} \lfloor (p-1-5j)/2 \rfloor + 1 \\ 2 \end{pmatrix},$$

and let

$$\widehat{L}(j) = \frac{j(5j-3)}{2} - \begin{pmatrix} \lfloor p-1-5j \rfloor / 2 + 2 \\ 2 \end{pmatrix}.$$

Then

$$L(2j) - L(2j-1) = \widehat{L}(2j) - \widehat{L}(2j-1) = p.$$

Lemma 2.2. *Let*

$$S_p = \{j \in \mathbb{Z} : 0 \leq \lfloor (p-1-5j)/2 \rfloor \leq p-1\}$$

and

$$\widehat{S}_p = \{j \in \mathbb{Z} : 0 \leq \lfloor (p-1-5j)/2 \rfloor + 1 \leq p-1\}.$$

We have

$$S_p = \{j \in \mathbb{Z} : -\lfloor p/5 \rfloor \leq j \leq \lfloor p/5 \rfloor\},$$

and

$$\widehat{S}_p = \begin{cases} \{j \in \mathbb{Z} : -\lfloor p/5 \rfloor + 1 \leq j \leq \lfloor p/5 \rfloor\} & \text{if } p \equiv 1 \pmod{5}, \\ \{j \in \mathbb{Z} : -\lfloor p/5 \rfloor \leq j \leq \lfloor p/5 \rfloor\} & \text{if } p \equiv 2, 3 \pmod{5}, \\ \{j \in \mathbb{Z} : -\lfloor p/5 \rfloor \leq j \leq \lfloor p/5 \rfloor + 1\} & \text{if } p \equiv 4 \pmod{5}. \end{cases}$$

These two lemmas above can be verified directly, so we omit the details here.

Proof of Theorem 1.1. By (2.1), we have

$$\mathcal{F}_{p+1}(q) = \sum_{j=-\infty}^{\infty} (-1)^j q^{j(5j+1)/2} \left[\lfloor (p-5j)/2 \rfloor \right]_q.$$

Observe that

$$\begin{bmatrix} p \\ k \end{bmatrix}_q \equiv \begin{cases} 1 \pmod{[p]_q} & \text{if } k = 1 \text{ or } p, \\ 0 \pmod{[p]_q} & \text{if } 1 \leq k \leq p-1. \end{cases}$$

Then it follows that

$$\mathcal{F}_{p+1}(q) \equiv \sum_{\lfloor (p-5j)/2 \rfloor = 0 \text{ or } p} (-1)^j q^{j(5j+1)/2} \pmod{[p]_q}.$$

It is easy to check that

$$\{j : \lfloor (p-5j)/2 \rfloor = 0 \text{ or } p\} = \begin{cases} \{(p-1)/5\} & \text{if } p \equiv 1 \pmod{5}, \\ \{-(p+1)/5\} & \text{if } p \equiv 4 \pmod{5}, \\ \emptyset & \text{if } p \equiv 2, 3 \pmod{5}. \end{cases}$$

Thus

$$\mathcal{F}_{p+1}(q) \equiv \sum_{\lfloor (p-5j)/2 \rfloor = 0 \text{ or } p} (-1)^j q^{j(5j+1)/2} \equiv \begin{cases} 1 \pmod{[p]_q} & \text{if } p \equiv 1, 4 \pmod{5}, \\ 0 \pmod{[p]_q} & \text{if } p \equiv 2, 3 \pmod{5}. \end{cases}$$

This concludes the proof of (1.4).

Also, applying (2.1) and Lemma 2.2, we deduce that

$$\begin{aligned} \mathcal{F}_p(q) &= \sum_{0 \leq \lfloor (p-1-5j)/2 \rfloor \leq p-1} (-1)^j q^{j(5j+1)/2} \left[\lfloor (p-1-5j)/2 \rfloor \right]_q \\ &= \sum_{j=-\lfloor p/5 \rfloor}^{\lfloor p/5 \rfloor} (-1)^j q^{j(5j+1)/2} \prod_{k=1}^{\lfloor (p-1-5j)/2 \rfloor} \frac{[p-k]_q}{[k]_q} \\ &= \sum_{j=-\lfloor p/5 \rfloor}^{\lfloor p/5 \rfloor} (-1)^j q^{j(5j+1)/2} \prod_{k=1}^{\lfloor (p-1-5j)/2 \rfloor} \frac{[p]_q - [k]_q}{q^k [k]_q} \\ &\equiv \sum_{j=-\lfloor p/5 \rfloor}^{\lfloor p/5 \rfloor} (-1)^{j+\lfloor (p-1-5j)/2 \rfloor} q^{L(j)} \pmod{[p]_q}. \end{aligned}$$

Suppose that $p \equiv 1 \pmod{5}$. Noting that $(p-1)/5$ is even and

$$(2j-1) + \lfloor (p-1-5(2j-1))/2 \rfloor = 2j + \lfloor (p-1-5 \cdot 2j)/2 \rfloor + 1 = (p-1)/2 - 3j + 1,$$

we have

$$\begin{aligned}
\mathcal{F}_p(q) &\equiv (-1)^{-(p-1)/5} q^{L(-(p-1)/5)} + \sum_{j=-(p-6)/5}^{(p-1)/5} (-1)^{j+\lfloor (p-1-5j)/2 \rfloor} q^{L(j)} \\
&= q^{(p-1)(p-2)/10 - p(p-1)/2} + \sum_{j=-(p-11)/10}^{(p-1)/10} (-1)^{(p-1)/2-3j} (q^{L(2j)} - q^{L(2j-1)}) \\
&\equiv q^{(4p+1)/5} \pmod{[p]_q}.
\end{aligned}$$

where Lemma 2.1 is applied in the last step. Similarly we obtain that

$$\mathcal{F}_p \equiv \begin{cases} (-1)^{(p-2)/5} q^{L((p-2)/5)} \equiv -q^{(2p+1)/5} \pmod{[p]_q} & \text{if } p \equiv 2 \pmod{5}, \\ (-1)^{-(p-3)/5} q^{L(-(p-3)/5)} \equiv -q^{(3p+1)/5} \pmod{[p]_q} & \text{if } p \equiv 3 \pmod{5}, \\ (-1)^{(p-4)/5} q^{L((p-4)/5)} \equiv q^{(p+1)/5} \pmod{[p]_q} & \text{if } p \equiv 4 \pmod{5}. \end{cases}$$

□

Proof of Theorem 1.2. In view of (2.2),

$$\widehat{\mathcal{F}}_p(q) = \sum_{j=-\infty}^{\infty} (-1)^j q^{j(5j-3)/2} \left[\lfloor (p-5j)/2 \rfloor + 1 \right]_q.$$

And we have

$$\{j : \lfloor (p-5j)/2 \rfloor + 1 = 0 \text{ or } p\} = \begin{cases} \{-(p-1)/5\} & \text{if } p \equiv 1 \pmod{5}, \\ \{-(p-2)/5\} & \text{if } p \equiv 2 \pmod{5}, \\ \{(p+2)/5\} & \text{if } p \equiv 3 \pmod{5}, \\ \{(p+1)/5\} & \text{if } p \equiv 4 \pmod{5}. \end{cases}$$

Therefore

$$\widehat{\mathcal{F}}_p(q) \equiv \begin{cases} (-1)^{-(p-1)/5} q^{(p+2)(p-1)/10} \equiv q^{(p-1)/5} \pmod{[p]_q} & \text{if } p \equiv 1 \pmod{5}, \\ (-1)^{-(p-2)/5} q^{(p+1)(p-2)/10} \equiv -q^{(3p-1)/5} \pmod{[p]_q} & \text{if } p \equiv 2 \pmod{5}, \\ (-1)^{(p+2)/5} q^{(p+2)(p-1)/10} \equiv -q^{(2p-1)/5} \pmod{[p]_q} & \text{if } p \equiv 3 \pmod{5}, \\ (-1)^{(p+1)/5} q^{(p+1)(p-2)/10} \equiv q^{(4p-1)/5} \pmod{[p]_q} & \text{if } p \equiv 4 \pmod{5}. \end{cases}$$

Now let us turn to the proof of (1.7). From (2.2), it follows that

$$\begin{aligned}
\widehat{\mathcal{F}}_{p-1}(q) &= \sum_{j=-\infty}^{\infty} (-1)^j q^{j(5j-3)/2} \left[\lfloor (p-1-5j)/2 \rfloor + 1 \right]_q \\
&= \sum_{0 \leq \lfloor (p-1-5j)/2 \rfloor + 1 \leq p-1}^{\infty} (-1)^j q^{j(5j-3)/2} \left[\lfloor (p-1-5j)/2 \rfloor + 1 \right]_q.
\end{aligned}$$

If $p \equiv 1 \pmod{5}$, then by Lemmas 2.1 and 2.2, we have

$$\begin{aligned}
\widehat{\mathcal{F}}_{p-1}(q) &= \sum_{j=-(p-1)/5+1}^{(p-1)/5} (-1)^j q^{j(5j-3)/2} \left[\left\lfloor \frac{p-1-5j}{2} \right\rfloor + 1 \right]_q \\
&\equiv \sum_{j=-(p-6)/5}^{(p-1)/5} (-1)^{j+\lfloor (p-1-5j)/2 \rfloor + 1} q^{\widehat{L}(j)} \\
&= \sum_{j=-(p-11)/10}^{(p-1)/10} (-1)^{(p-1)/2-3j+1} (q^{\widehat{L}(2j)} - q^{\widehat{L}(2j-1)}) \\
&\equiv 0 \pmod{[p]_q}.
\end{aligned}$$

Similarly when $p \equiv 4 \pmod{5}$,

$$\widehat{\mathcal{F}}_{p-1}(q) \equiv \sum_{j=-(p-4)/5}^{(p+1)/5} (-1)^{j+\lfloor (p-1-5j)/2 \rfloor + 1} q^{\widehat{L}(j)} \equiv 0 \pmod{[p]_q}.$$

Finally, suppose that $p \equiv 2$ or $3 \pmod{5}$. Then

$$\begin{aligned}
&\widehat{\mathcal{F}}_{p-1}(q) \\
&\equiv \sum_{j=-\lfloor p/5 \rfloor}^{\lfloor p/5 \rfloor} (-1)^{j+\lfloor (p-1-5j)/2 \rfloor + 1} q^{\widehat{L}(j)} \\
&\equiv \begin{cases} (-1)^{(p-2)/5+1} q^{\widehat{L}((p-2)/5)} = q^{p(p-7)/10} \equiv 1 \pmod{[p]_q} & \text{if } p \equiv 2 \pmod{5}, \\ (-1)^{-(p-3)/5+(p-1)} q^{\widehat{L}(-(p-3)/5)} = q^{p(1-2p)/5} \equiv 1 \pmod{[p]_q} & \text{if } p \equiv 3 \pmod{5}. \end{cases}
\end{aligned}$$

All are done. $\square$

3. q -PELL NUMBER

To prove Theorem 1.3, we need the similar identities as (2.1) and (2.2) for $\mathcal{P}_n(q)$ and $\widehat{\mathcal{P}}_n(q)$ respectively. Fortunately, such identities have been established by Santos and Sills in [8]. Let

$$T_1(n, m, q) = \sum_{j=0}^n (-q)^j \begin{bmatrix} n \\ j \end{bmatrix}_{q^2} \begin{bmatrix} 2n-2j \\ n-m-j \end{bmatrix}_q.$$

Lemma 3.1 (Santos and Sills, [8]). *Let n be a non-negative integer. Then*

$$\begin{aligned}
\mathcal{P}_{n+1}(q) &= \sum_{j=0}^n \sum_{k=0}^j q^{(j^2+j+k^2-k)/2} \begin{bmatrix} j \\ k \end{bmatrix}_q \begin{bmatrix} n-k \\ j \end{bmatrix}_q \\
&= \sum_{j=-\infty}^{\infty} (-1)^j q^{2j^2} T_1(n+1, 4j+1, \sqrt{q}), \tag{3.1}
\end{aligned}$$

and

$$\begin{aligned}\widehat{\mathcal{P}}_{n+1}(q) &= \sum_{j=0}^n \sum_{k=0}^j q^{(j^2+j+k^2+k)/2} \begin{bmatrix} j \\ k \end{bmatrix}_q \begin{bmatrix} n-k \\ j \end{bmatrix}_q \\ &= \sum_{j=-\infty}^{\infty} (-1)^j q^{2j^2+j} T_1(n+1, 4j+1, \sqrt{q}).\end{aligned}\quad (3.2)$$

Proof of Theorem 1.3. For arbitrary two polynomials $f(q)$ and $g(q)$, it is easy to see that $f(q^2) \mid g(q^2)$ implies $f(q) \mid g(q)$. Indeed, if

$$g(q^2)/f(q^2) = \sum_{k=0}^n a_k q^k,$$

then

$$\sum_{k=0}^n a_k (-q)^k = g((-q)^2)/f((-q)^2) = g(q^2)/f(q^2) = \sum_{k=0}^n a_k q^k,$$

whence $a_k = 0$ for each odd k . Thus

$$g(q)/f(q) = \sum_{0 \leq k \leq n/2} a_{2k} q^k.$$

Now it suffices to show that

$$q^{(p^2-1)/4} \mathcal{P}_p(q^2) \equiv \left(\frac{2}{p}\right) \pmod{[p]_{q^2}}$$

and

$$\widehat{\mathcal{P}}_p(q^2) \equiv \left(\frac{2}{p}\right) q^{(p^2-1)/4} \pmod{[p]_{q^2}}.$$

By (3.1), we have

$$\begin{aligned}\mathcal{P}_p(q^2) &= \sum_{j=-\infty}^{\infty} (-1)^j q^{4j^2} T_1(n+1, 4j+1, q) \\ &= \sum_{j=-\infty}^{\infty} (-1)^j q^{4j^2} \sum_{k=0}^p (-q)^k \begin{bmatrix} p \\ k \end{bmatrix}_{q^2} \begin{bmatrix} 2p-2k \\ p-4j-k-1 \end{bmatrix}_q \\ &\equiv \sum_{j=-\infty}^{\infty} (-1)^j q^{4j^2} \begin{bmatrix} 2p \\ p-4j-1 \end{bmatrix}_q \pmod{[p]_{q^2}}.\end{aligned}$$

Since

$$\begin{bmatrix} 2p \\ k \end{bmatrix}_q \equiv 0 \pmod{[p]_{q^2}}$$

when $p \nmid k$, we have

$$\begin{aligned} \mathcal{P}_p(q^2) &\equiv \begin{cases} (-1)^{(p-1)/4} q^{(p-1)^2/4} & \text{if } p \equiv 1 \pmod{4} \\ (-1)^{-(p+1)/4} q^{(p+1)^2/4} & \text{if } p \equiv 3 \pmod{4} \end{cases} \\ &\equiv \left(\frac{2}{p}\right) q^{-(p^2-1)/4} \pmod{[p]_{q^2}}. \end{aligned}$$

The proof of (1.10) is very similar. From (3.2), we deduce that

$$\begin{aligned} \widehat{\mathcal{P}}_p(q^2) &= \sum_{j=-\infty}^{\infty} (-1)^j q^{4j^2+2j} \sum_{k=0}^p (-q)^k \begin{bmatrix} p \\ k \end{bmatrix}_{q^2} \begin{bmatrix} 2p-2k \\ p-4j-k-1 \end{bmatrix}_q \\ &\equiv \sum_{j=-\infty}^{\infty} (-1)^j q^{4j^2+2j} \begin{bmatrix} 2p \\ p-4j-1 \end{bmatrix}_q \\ &\equiv \begin{cases} (-1)^{(p-1)/4} q^{(p-1)^2/4+(p-1)/2} & \text{if } p \equiv 1 \pmod{4} \\ (-1)^{-(p+1)/4} q^{(p+1)^2/4-(p+1)/2} & \text{if } p \equiv 3 \pmod{4} \end{cases} \\ &= \left(\frac{2}{p}\right) q^{(p^2-1)/4} \pmod{[p]_{q^2}}. \end{aligned}$$

Thus the proofs of (1.9) and (1.10) are completed.

We need the following lemma in the proof of (1.11).

Lemma 3.2.

$$\begin{bmatrix} 2p+2 \\ p \end{bmatrix}_q \equiv 1 + q^p \pmod{[p]_{q^2}}$$

for any odd prime p .

Proof. Write

$$\begin{bmatrix} 2p+2 \\ p \end{bmatrix}_q = \frac{(1-q^{2p+2})(1-q^{2p+1})(1-q^{2p}) \prod_{j=1}^{p-1} (1-q^{2p-j})}{(1-q^{p+2})(1-q^{p+1})(1-q^p) \prod_{j=1}^{p-1} (1-q^j)}.$$

Since

$$\frac{1-q^{2p-j}}{1-q^j} \equiv \frac{q^j - q^{2p}}{q^j(1-q^j)} = \frac{(1-q^{2p}) - (1-q^j)}{q^j(1-q^j)} \equiv -q^{-j} \pmod{[p]_{q^2}},$$

we have

$$\begin{bmatrix} 2p+2 \\ p \end{bmatrix}_q \equiv (-1)^{p-1} q^{-\binom{p}{2}} \frac{(1-q^{2p+2})(1-q^{2p+1})(1-q^{2p})}{(1-q^{p+2})(1-q^{p+1})(1-q^p)} \pmod{[p]_{q^2}}.$$

And noting that

$$q^{-\binom{p}{2}} \frac{(1 - q^{2p+2})(1 - q^{2p+1})}{(1 - q^{p+2})(1 - q^{p+1})} \equiv 1 \pmod{[p]_q},$$

it follows that

$$q^{-\binom{p}{2}} \frac{(1 - q^{2p+2})(1 - q^{2p+1})(1 - q^{2p})}{(1 - q^{p+2})(1 - q^{p+1})(1 - q^p)} \equiv 1 + q^p \pmod{(1 + q^p)[p]_q = (1 + q)[p]_{q^2}}.$$

We are done. $\square$

From the definition of T_1 , we have

$$\begin{aligned} & T_1(p+1, 4j+1, q) + qT_1(p, 4j+1, q) \\ &= \sum_{k=0}^{p+1} (-q)^k \begin{bmatrix} p+1 \\ k \end{bmatrix}_{q^2} \begin{bmatrix} 2p+2-2k \\ p-4j-k \end{bmatrix}_q - \sum_{k=0}^p (-q)^{k+1} \begin{bmatrix} p \\ k \end{bmatrix}_{q^2} \begin{bmatrix} 2p-2k \\ p-4j-k-1 \end{bmatrix}_q \\ &= \begin{bmatrix} 2p+2 \\ p-4j \end{bmatrix}_q + \sum_{k=0}^p (-q)^{k+1} \left(\begin{bmatrix} p+1 \\ k+1 \end{bmatrix}_{q^2} - \begin{bmatrix} p \\ k \end{bmatrix}_{q^2} \right) \begin{bmatrix} 2p-2k \\ p-4j-k-1 \end{bmatrix}_q \\ &= \begin{bmatrix} 2p+2 \\ p-4j \end{bmatrix}_q + \sum_{k=0}^p (-q)^{k+1} q^{2(k+1)} \begin{bmatrix} p \\ k+1 \end{bmatrix}_{q^2} \begin{bmatrix} 2p-2k \\ p-4j-k-1 \end{bmatrix}_q \\ &\equiv \begin{bmatrix} 2p+2 \\ p-4j \end{bmatrix}_q + (-q)^p \begin{bmatrix} 2 \\ -4j \end{bmatrix}_q \pmod{[p]_{q^2}}. \end{aligned}$$

Thus

$$\begin{aligned} \mathcal{P}_{p+1}(q^2) + q\mathcal{P}_p(q^2) &= \sum_{j=-\infty}^{\infty} (-1)^j q^{4j^2} (T_1(p+1, 4j+1, q) - qT_1(p, 4j+1, q)) \\ &\equiv \sum_{j=-\infty}^{\infty} (-1)^j q^{4j^2} \begin{bmatrix} 2p+2 \\ p-4j \end{bmatrix}_q + \sum_{j=-\infty}^{\infty} (-1)^{p-j} q^{4j^2+p} \begin{bmatrix} 2 \\ -4j \end{bmatrix}_q \\ &\equiv \sum_{j=-\infty}^{\infty} (-1)^j q^{4j^2} \begin{bmatrix} 2p+2 \\ p-4j \end{bmatrix}_q - q^p \pmod{[p]_{q^2}}. \end{aligned}$$

Note that

$$\begin{bmatrix} 2p+2 \\ k \end{bmatrix}_q \not\equiv 0 \pmod{[p]_{q^2}}$$

only if $k \in \{0, 1, 2, p, p+1, p+2, 2p, 2p+1, 2p+2\}$. Hence we have

$$\begin{aligned}
 & \sum_{j=-\infty}^{\infty} (-1)^j q^{4j^2} \begin{bmatrix} 2p+2 \\ p-4j \end{bmatrix}_q \\
 & \equiv \begin{bmatrix} 2p+2 \\ p \end{bmatrix}_q + \begin{cases} (-1)^{(p-1)/4} q^{(p-1)^2/4} \begin{bmatrix} 2p+2 \\ 1 \end{bmatrix}_q & \text{if } p \equiv 1 \pmod{4} \\
 (-1)^{-(p+1)/4} q^{(p+1)^2/4} \begin{bmatrix} 2p+2 \\ 2p+1 \end{bmatrix}_q & \text{if } p \equiv 3 \pmod{4} \end{cases} \\
 & \equiv 1 + q^p + \left(\frac{2}{p}\right) q^{-(p^2-1)/4} (1+q) \\
 & \equiv 1 + q^p + (1+q) \mathcal{P}_p(q^2) \pmod{[p]_{q^2}}.
 \end{aligned}$$

This concludes that

$$\mathcal{P}_{p+1}(q^2) \equiv 1 + \mathcal{P}_p(q^2) \pmod{[p]_{q^2}}.$$

Similarly, we have

$$\begin{aligned}
 \widehat{\mathcal{P}}_{p+1}(q^2) + q \widehat{\mathcal{P}}_p(q^2) &= \sum_{j=-\infty}^{\infty} (-1)^j q^{4j^2+2j} (T_1(p+1, 4j+1, q) - q T_1(p, 4j+1, q)) \\
 &\equiv \sum_{j=-\infty}^{\infty} (-1)^j q^{4j^2+2j} \begin{bmatrix} 2p+2 \\ p-4j \end{bmatrix}_q - q^p \\
 &\equiv \begin{bmatrix} 2p+2 \\ p \end{bmatrix}_q + \left(\frac{2}{p}\right) q^{(p^2-1)/4} (1+q) - q^p \\
 &\equiv 1 + q^p + (1+q) \widehat{\mathcal{P}}_p(q^2) - q^p \pmod{[p]_{q^2}},
 \end{aligned}$$

whence

$$\widehat{\mathcal{P}}_{p+1}(q^2) \equiv 1 + \widehat{\mathcal{P}}_p(q^2) \pmod{[p]_{q^2}}.$$

□

Acknowledgment. I thank my advisor, Zhi-Wei Sun, for his help on this paper.

REFERENCES

- [1] G. E. Andrews, *A polynomial identity which implies the Rogers-Ramanujan identities*, Scripta Math. **28** (1970), 297-305.
- [2] G. E. Andrews, *Fibonacci numbers and the Rogers-Ramanujan identities*, Fibonacci Quarterly **42** (2004), 3-19.
- [3] A. Berkovich and P. Paule, *Variants of the Andrews-Gordon identities*, Ramanujan J. **5** (2001), 391-404.
- [4] L. Carlitz, *Fibonacci notes. III: q -Fibonacci numbers*, Fibonacci Quarterly **12** (1974), 317-322.

- [5] J. Cigler, *q-Fibonacci polynomials*, Fibonacci Quarterly **41** (2003), 31-40.
- [6] K. Garrett, M. Ismail and D. Stanton, *Variants of the Rogers-Ramanujan identities*, Adv. Appl. Math. **23** (1999), 274-299.
- [7] P. Ribenboim, *The Book of Prime Number Records*, 2nd ed., Springer-Verlag, New York, 1989, pp. 44-50.
- [8] J. Santos and A. V. Sills, *q-Pell sequences and two identities of V. A. Lebesgue*, Discrete Math. **257** (2002), 125-142.
- [9] I. Schur, *Gesammelte Abhandlungen*, Vol. 2, Springer-Verlag, Berlin, 1973, pp. 117-136.
- [10] Z. H. Sun, *Values of Lucas sequences modulo primes*, Rocky Mountain J. Math. **33** (2003), 1123-1145.
- [11] Z. H. Sun and Z. W. Sun, *Fibonacci numbers and Fermat's last theorem*, Acta Arith. **60** (1992), 371-388.
- [12] H. C. Williams, *A note on the Fibonacci quotient $F_{p-\epsilon}/p$* , Can. Math. Bull. **25** (1982), 366-370.

DEPARTMENT OF MATHEMATICS, NANJING UNIVERSITY, NANJING 210093, PEOPLE'S REPUBLIC OF CHINA

E-mail address: haopan79@yahoo.com.cn